

COMODULES FOR SOME SIMPLE $\mathcal{O}$ -FORMS OF $\mathbb{G}_m$

N. E. CSIMA AND R. E. KOTTWITZ

ABSTRACT. This paper gives a rather concrete description of the category $\text{Rep}(G)$ for certain flat commutative affine group schemes G over a discrete valuation ring such that the general fiber of G is the multiplicative group.

Tannakian theory allows one to understand an affine group scheme G over a commutative base ring A in terms of the category $\text{Rep}(G)$ of G -modules, by which is meant comodules for the Hopf algebra corresponding to G . The theory is especially well-developed [Sa] in the case that A is a field, and some parts of the theory still work well over more general rings A , say discrete valuation rings (see [Sa, We]).

When A is a field of characteristic zero and G is connected reductive, the category $\text{Rep}(G)$ is very well understood. However, with the exception of groups as simple as the multiplicative and additive groups, little seems to be known about what $\text{Rep}(G)$ looks like concretely when A is no longer assumed to be a field, even in the most favorable case in which A is a discrete valuation ring and G is a flat affine group scheme over A with connected reductive general fiber.

The modest goal of this paper is to give a concrete description of $\text{Rep}(G)$ for certain flat group schemes G over a discrete valuation ring $\mathcal{O}$ such that the general fiber of G is $\mathbb{G}_m$.

Choose a generator π of the maximal ideal of $\mathcal{O}$ and write F for the field of fractions of $\mathcal{O}$. For any non-negative integer k , the construction of 1.1, when applied to $f = \pi^k$, yields a commutative flat affine group scheme G_k over $\mathcal{O}$ whose general fiber is $\mathbb{G}_m$. The $\mathcal{O}$ -points of G_k are given by

$$G_k(\mathcal{O}) = \{t \in \mathcal{O}^\times : t \equiv 1 \pmod{\pi^k}\}.$$

These form a projective system

$$\cdots \rightarrow G_2 \rightarrow G_1 \rightarrow G_0 = \mathbb{G}_m$$

in an obvious way, and we may form the projective limit $G_\infty := \text{proj lim } G_k$. The Hopf algebra S_k corresponding to G_k can be described explicitly (see 1.1 and 1.2). The Hopf algebra S_∞ corresponding to G_∞ is

$$\text{inj lim } S_k = \left\{ \sum_{i \in \mathbb{Z}} x_i T^i \in F[T, T^{-1}] : \sum_{i \in \mathbb{Z}} x_i \in \mathcal{O} \right\}.$$

The categories $\text{Rep}(G_\infty)$ and $\text{Rep}(G_k)$ can be described very concretely. Indeed, $\text{Rep}(G_\infty)$ consists of the category of $\mathcal{O}$ -modules M equipped with a $\mathbb{Z}$ -grading on $F \otimes_{\mathcal{O}} M$ (see 2.3, where a much more general result is proved). As for $\text{Rep}(G_k)$, we proceed in two steps.

First, the full subcategory of $\text{Rep}(G_k)$ consisting of those G_k -modules that are flat as $\mathcal{O}$ -modules is equivalent (see Theorem 1.3.1) to the category of pairs (V, M)

1991 *Mathematics Subject Classification*. Primary 16W30; Secondary 14L17.
Partially supported by NSF Grant DMS-0245639.

consisting of a $\mathbb{Z}$ -graded F -vector space V and an *admissible* $\mathcal{O}$ -submodule M of V , where admissible means that the canonical map $F \otimes_{\mathcal{O}} M \rightarrow V$ is an isomorphism and $C_n M \subset M$ for all $n \geq 0$, where $C_n : V \rightarrow V$ is the graded linear map given by multiplication by $\pi^{kn} \binom{i}{n}$ on the i -th graded piece of V . The G_k -module corresponding to (V, M) is M , equipped with the obvious comultiplication.

Second, any G_k -module (see 1.4) is obtained as the cokernel of some injective homomorphism $M_1 \rightarrow M_0$ coming from a morphism $(V_1, M_1) \rightarrow (V_0, M_0)$ of pairs of the type just described.

When $\mathcal{O}$ is a $\mathbb{Q}$ -algebra, the situation is even simpler: M is an admissible $\mathcal{O}$ -submodule of the graded vector space V if and only if $C_1 M \subset M$ and $F \otimes_{\mathcal{O}} M \cong V$. Moreover, in case $\mathcal{O}$ is the formal power series ring $\mathbb{C}[[\varepsilon]]$, there is an interesting connection with affine Springer fibers (see 1.5).

1. A DESCRIPTION OF $\text{Rep}(G)_f$ FOR CERTAIN GROUP SCHEMES G

Throughout this section we consider a commutative ring A and a nonzerodivisor $f \in A$. Thus the canonical homomorphism $A \rightarrow A_f$ is injective, where A_f denotes the localization of A with respect to the multiplicative subset $\{f^n : n \geq 0\}$. For the rest of this section we denote A_f by B and use the canonical injection $A \hookrightarrow B$ to identify A with a subring of B .

1.1. The group scheme G over A . We are now going to define a commutative affine group scheme G , flat and finitely presented over A . There will be a canonical homomorphism $G \rightarrow \mathbb{G}_m$ that becomes an isomorphism after extending scalars from A to B .

We begin by specifying the functor of points for G . For any commutative A -algebra R we put

$$\begin{aligned} G(R) &:= \{(t, x) \in R^\times \times R : t - 1 = fx\} \\ &= \{x \in R : 1 + fx \in R^\times\}. \end{aligned}$$

Then G is represented by the A -algebra

$$\begin{aligned} (1.1.1) \quad S &:= A[T, T^{-1}, X]/(T - 1 - fX) \\ &= A[X]_{1+fX}, \end{aligned}$$

which is clearly flat and finitely presented.

The multiplication on $G(R)$ is defined as $(t, x)(t', x') = (tt', x + x' + fxx')$. The identity element is $(1, 0)$ and the inverse of (t, x) is $(t^{-1}, -t^{-1}x)$.

There is a canonical homomorphism $\lambda : G \rightarrow \mathbb{G}_m$, given by $(t, x) \mapsto t$. When f is a nonzerodivisor in R , the homomorphism $\lambda : G(R) \rightarrow R^\times$ identifies $G(R)$ with $\ker[R^\times \rightarrow (R/fR)^\times]$, and when f is a unit in R , then $G(R) = R^\times$, showing that the homomorphism $\lambda : G \rightarrow \mathbb{G}_m$ becomes an isomorphism after extending scalars from A to B . Thus there is a canonical isomorphism $B \otimes_A S \cong B[T, T^{-1}]$.

Lemma 1.1.1. *Let M be an A -module on which f is a nonzerodivisor. Let F be any flat A -module. Then f is also a nonzerodivisor on $F \otimes_A M$.*

Proof. Tensor the injection $M \xrightarrow{f} M$ over A with F . □

Corollary 1.1.2. *The canonical homomorphism $S \rightarrow B \otimes_A S = B[T, T^{-1}]$ is injective, so that we may identify S with a subring of $B[T, T^{-1}]$.*

Proof. Just note that S is flat over A and f is a nonzerodivisor on A . Therefore f is a nonzerodivisor on $S \otimes_A A = S$, and this means that $S \rightarrow B \otimes_A S$ is injective. $\square$

1.2. Description of S as a subring of $B[T, T^{-1}]$. We have just identified S with a subring of $B[T, T^{-1}]$. It is obvious from (1.1.1) that S is the A -subalgebra of $B[T, T^{-1}]$ generated by $T, T^{-1}, \frac{T-1}{f}$. However there is a more useful description of S in terms of B -module maps

$$L_n : B[T, T^{-1}] \rightarrow B,$$

one for each non-negative integer n , defined by the formula

$$L_n\left(\sum_{i \in \mathbb{Z}} b_i T^i\right) = \sum_{i \in \mathbb{Z}} f^n \binom{i}{n} b_i.$$

Here $\binom{i}{n}$ is the binomial coefficient $i(i-1)\dots(i-n+1)/n!$, defined for all $i \in \mathbb{Z}$. When $n = 0$, we have $\binom{i}{n} = 1$ for all $i \in \mathbb{Z}$.

The following remarks may help in understanding the maps L_n . For any non-negative integer n , we have the divided-power differential operator

$$D^{[n]} : B[T, T^{-1}] \rightarrow B[T, T^{-1}]$$

defined by

$$(1.2.1) \quad D^{[n]}\left(\sum_{i \in \mathbb{Z}} b_i T^i\right) = \sum_{i \in \mathbb{Z}} \binom{i}{n} b_i T^{i-n}.$$

The Leibniz formula says that

$$(1.2.2) \quad D^{[n]}(gh) = \sum_{r=0}^n D^{[r]}(g) D^{[n-r]}(h).$$

For any $g \in B[T] \subset B[T, T^{-1}]$ the Taylor expansion of g at $T = 1$ reads

$$(1.2.3) \quad g = \sum_{n=0}^{\infty} (D^{[n]}g)(1) \cdot (T-1)^n,$$

the sum having only finitely many non-zero terms.

For any $g \in B[T, T^{-1}]$ we have $L_n(g) = f^n (D^{[n]}g)(1)$. It follows from (1.2.2) that for all $g, h \in B[T, T^{-1}]$

$$(1.2.4) \quad L_n(gh) = \sum_{r=0}^n L_r(g) L_{n-r}(h),$$

and for all $h \in B[T] \subset B[T, T^{-1}]$ it follows from (1.2.3) that

$$(1.2.5) \quad h = \sum_{n=0}^{\infty} L_n(h) \left(\frac{T-1}{f}\right)^n.$$

Now we are in a position to prove

Proposition 1.2.1. *The subring S of $B[T, T^{-1}]$ is equal to*

$$\{g \in B[T, T^{-1}] : \forall n \geq 0 \ L_n(g) \in A\}.$$

Proof. Write S' for $\{g \in B[T, T^{-1}] : \forall n \geq 0 \ L_n(g) \in A\}$. Obviously S' is an A -submodule of $B[T, T^{-1}]$, and it follows from (1.2.4) that S' is a subring of $B[T, T^{-1}]$. A simple calculation shows that $T, T^{-1}, (T-1)/f$ lie in S' , and as these three elements generate S as A -algebra, we conclude that $S \subset S'$.

Now let $g \in S'$. There exists an integer n large enough that $h := T^n g$ lies in the subring $B[T]$. Note that $h \in S'$. Equation (1.2.5) shows that $h \in S$, since $(T-1)/f \in S$ and $L_n(h) \in A$. Therefore $g = T^{-n}h \in S$. $\square$

Now let M be an A -module on which f is a nonzerodivisor, so that we may use the canonical A -module map $M \rightarrow B \otimes_A M$ (sending m to $1 \otimes m$) to identify M with an A -submodule of $N := B \otimes_A M$.

It follows from Lemma 1.1.1 that the canonical A -module map

$$S \otimes_A M \rightarrow B \otimes_A (S \otimes_A M) = B[T, T^{-1}] \otimes_B N$$

identifies $S \otimes_A M$ with an A -submodule of $B[T, T^{-1}] \otimes_B N$. We will now derive from the previous proposition a description of $S \otimes_A M$ inside $B[T, T^{-1}] \otimes_B N$. For this we will need the B -module maps $\mathbf{L}_n : B[T, T^{-1}] \otimes_B N \rightarrow N$ defined by

$$\mathbf{L}_n\left(\sum_{i \in \mathbb{Z}} T^i \otimes x_i\right) = \sum_{i \in \mathbb{Z}} f^n \binom{i}{n} x_i.$$

Here $x_i \in N$, all but finitely many being 0.

Lemma 1.2.2. *The A -submodule $S \otimes_A M$ of $B[T, T^{-1}] \otimes_B N$ is equal to*

$$\{x \in B[T, T^{-1}] \otimes_B N : \forall n \geq 0 \ \mathbf{L}_n(x) \in M\}.$$

Proof. From Proposition 1.2.1 we see that there is an exact sequence

$$0 \rightarrow S \rightarrow B[T, T^{-1}] \xrightarrow{L} \prod_{n \geq 0} B/A,$$

the n -th component of the map L being the composition

$$B[T, T^{-1}] \xrightarrow{L_n} B \twoheadrightarrow B/A.$$

In fact the map L takes values in $\oplus_{n \geq 0} B/A$. Indeed, for any $g \in B[T, T^{-1}]$ there exists an integer m large enough that $f^m g \in A[T, T^{-1}]$, and then $L_n(g) \in A$ for all $n \geq m$. Moreover L maps $B[T, T^{-1}]$ onto $\oplus_{n \geq 0} B/A$. Indeed, a simple calculation shows that for $b \in B$ and $m \geq 0$

$$L_n(b f^{-m} (T-1)^m) = \begin{cases} b & \text{if } m = n \\ 0 & \text{otherwise.} \end{cases}$$

(First check that $D^{[n]}((T-1)^m) = \binom{m}{n} (T-1)^{m-n}$, say by induction on m ; note that this formula is valid even if $n > m$, since $\binom{m}{n} = 0$ when $0 \leq m < n$.)

We now have a short exact sequence

$$0 \rightarrow S \rightarrow B[T, T^{-1}] \xrightarrow{L} \bigoplus_{n \geq 0} B/A \rightarrow 0$$

of A -modules. Tensoring with the A -module M , we obtain an exact sequence

$$(1.2.6) \quad S \otimes_A M \rightarrow B[T, T^{-1}] \otimes_A M \xrightarrow{L \otimes id_M} \left(\bigoplus_{n \geq 0} B/A\right) \otimes_A M \rightarrow 0.$$

Now

$$B[T, T^{-1}] \otimes_A M = B[T, T^{-1}] \otimes_B B \otimes_A M = B[T, T^{-1}] \otimes_B N$$

and

$$\left(\bigoplus_{n \geq 0} B/A \right) \otimes_A M = \bigoplus_{n \geq 0} N/M.$$

With these identifications (and recalling that $S \otimes_A M \rightarrow B[T, T^{-1}] \otimes_B N$ is injective), we see that (1.2.6) describes $S \otimes_A M$ as the subset of $B[T, T^{-1}] \otimes_B N$ consisting of elements x such that $\mathbf{L}_n(x) \in M$ for all $n \geq 0$, and this completes the proof. $\square$

1.3. Comodules for S . Since G is an affine group scheme over A , the A -algebra S is actually a commutative Hopf algebra, and we can consider $\text{Rep}(G)$, the category of S -comodules. We denote by $\text{Rep}(G)_f$ the full subcategory of $\text{Rep}(G)$ consisting of S -comodules M such that f is a nonzerodivisor on the A -module underlying M . Our next goal is to give a concrete description of $\text{Rep}(G)_f$.

In order to do so, we need one more construction. Let $N = \bigoplus_{i \in \mathbb{Z}} N_i$ be a $\mathbb{Z}$ -graded B -module. For each non-negative integer n we define an endomorphism $C_n : N \rightarrow N$ of the graded B -module N by requiring that C_n be given by multiplication by $f^n \binom{i}{n}$ on N_i . Thus

$$C_n \left(\sum_{i \in \mathbb{Z}} x_i \right) = \sum_{i \in \mathbb{Z}} f^n \binom{i}{n} x_i.$$

Here $x_i \in N_i$, all but finitely many being 0.

Let $\mathcal{C}$ be the category whose objects are pairs (N, M) , N being a $\mathbb{Z}$ -graded B -module, and M being an A -submodule of N such that the natural map $B \otimes_A M \rightarrow N$ is an isomorphism and such that $C_n M \subset M$ for all $n \geq 0$. A morphism $(N, M) \rightarrow (N', M')$ is a homomorphism $\phi : N \rightarrow N'$ of graded B -modules such that $\phi M \subset M'$.

We now define a functor $F : \text{Rep}(G)_f \rightarrow \mathcal{C}$. Let M be an object of $\text{Rep}(G)_f$. Then $N := B \otimes_A M$ is a comodule for $B \otimes_A S = B[T, T^{-1}]$. It is known (see [SGA3], Exposé 1) that the category of $B[T, T^{-1}]$ -comodules is equivalent to the category of $\mathbb{Z}$ -graded B -modules. Thus N has a $\mathbb{Z}$ -grading $N = \bigoplus_{i \in \mathbb{Z}} N_i$, and the comultiplication $\Delta_N : N \rightarrow B[T, T^{-1}] \otimes_B N$ is given by $\sum_{i \in \mathbb{Z}} x_i \mapsto \sum_{i \in \mathbb{Z}} T^i \otimes x_i$. Since f is a nonzerodivisor on M , the canonical map $M \rightarrow B \otimes_A M = N$ identifies M with an A -submodule of N .

We define our functor F by $FM := (N, M)$. For this to make sense we must check that $C_n M \subset M$ for all $n \geq 0$. Let $m \in M$, and write $m = \sum_{i \in \mathbb{Z}} x_i$ in $\bigoplus_{i \in \mathbb{Z}} N_i = N$. Since the comodule N was obtained from M by extension of scalars, the element $x = \Delta_N m = \sum_{i \in \mathbb{Z}} T^i \otimes x_i \in B[T, T^{-1}] \otimes_B N$ lies in the image of $S \otimes_A M \rightarrow B[T, T^{-1}] \otimes_B N$. Lemma 1.2.2 then implies that $\mathbf{L}_n(x) = \sum_{i \in \mathbb{Z}} f^n \binom{i}{n} x_i = C_n(m)$ lies in M , as desired.

Theorem 1.3.1. *The functor $F : \text{Rep}(G)_f \rightarrow \mathcal{C}$ is an equivalence of categories.*

Proof. Let us first show that F is essentially surjective. Let (N, M) be an object in $\mathcal{C}$. We are going to use the comultiplication $\Delta_N : N \rightarrow B[T, T^{-1}] \otimes_B N$ to turn M into an S -comodule.

Since M is an A -submodule of N , it is clear that f is a nonzerodivisor on M . As we have seen before, it follows that f is a nonzerodivisor on $S \otimes_A M$ and hence that the natural map $S \otimes_A M \rightarrow B \otimes_A (S \otimes_A M) = B[T, T^{-1}] \otimes_B N$ identifies $S \otimes_A M$ with an A -submodule of $B[T, T^{-1}] \otimes_B N$.

Using Lemma 1.2.2, we see that our assumption that $C_n M \subset M$ for all $n \geq 0$ is simply the statement that $\Delta_N M \subset S \otimes_A M$. In other words there exists a unique A -module map $\Delta_M : M \rightarrow S \otimes_A M$ such that Δ_M yields Δ_N after extending scalars from A to B .

We claim that Δ_M makes M into an S -comodule. For this we must check the commutativity of two diagrams, and this follows from the commutativity of these diagrams after extending scalars from A to B , once one notes that for any two A -modules M_1, M_2 on which f is a nonzerodivisor

$$(1.3.1) \quad \text{Hom}_A(M_1, M_2) = \{\phi \in \text{Hom}_B(B \otimes_A M_1, B \otimes_A M_2) : \phi(M_1) \subset M_2\}.$$

Here of course we are identifying M_1, M_2 with A -submodules of $B \otimes_A M_1, B \otimes_A M_2$ respectively. (At one point we need that f is a nonzerodivisor on $S \otimes_A S \otimes_A M$, which is true since $S \otimes_A S$ is flat over A .)

As F takes M to (N, M) , we are done with essential surjectivity. It is easy to see that F is fully faithful; this too uses (1.3.1). $\square$

1.4. Principal ideal domains A . One defect of the theorem we just proved is that it only describes those G -modules on which f is a nonzerodivisor. When A is a principal ideal domain, as we assume for the rest of this subsection, we can do better. Now f is simply any non-zero element of A . As a consequence of our theorem we obtain an equivalence of categories between the category $\text{Rep}(G)_{\text{flat}}$ of G -modules M such that M is flat as A -module and the full subcategory of $\mathcal{C}$ consisting of pairs (N, M) for which M is a flat A -module (in which case $N \cong B \otimes_A M$ is necessarily a flat B -module).

The next lemma is a variant of [Se, Prop. 3].

Lemma 1.4.1. *Let A be a principal ideal domain, let C be a flat A -coalgebra, and let E be a C -comodule. Then there exists a short exact sequence of C -comodules*

$$0 \rightarrow F_1 \rightarrow F_0 \rightarrow E \rightarrow 0$$

in which F_0, F_1 are flat as A -modules.

Proof. We imitate Serre's proof. Recall (see 1.2 in *loc. cit.*) that for any A -module M the map $\Delta \otimes id_M : C \otimes_A M \rightarrow C \otimes_A C \otimes_A M$ (Δ being the comultiplication for C) gives $C \otimes_A M$ the structure of C -comodule, and that (see 1.4 in *loc. cit.*) the comultiplication map $\Delta_E : E \rightarrow C \otimes_A E$ is an injective comodule map when $C \otimes_A E$ is given the comodule structure just described. We use Δ_E to identify E with a subcomodule of $C \otimes_A E$.

Now choose a surjective A -linear map $p : F \rightarrow E$, where F is a free A -module. Let F_0 denote the preimage of E under the surjective comodule map $id \otimes p : C \otimes_A F \rightarrow C \otimes_A E$. Since F_0 is the kernel of

$$C \otimes_A F \rightarrow C \otimes_A E \rightarrow (C \otimes_A E)/E,$$

it is a subcomodule of $C \otimes_A F$, and $id \otimes p$ restricts to a surjective comodule map $F_0 \rightarrow E$, whose kernel we denote by F_1 . Since C and F are flat, so too are $C \otimes_A F$, F_0 , and F_1 , and we are done. We used that for principal ideal domains, a module is flat if and only if it is torsion-free, and being torsion-free is a property that is inherited by submodules. $\square$

Returning to our Hopf algebra S , we see that any G -module E has a resolution $0 \rightarrow F_1 \rightarrow F_0 \rightarrow E \rightarrow 0$ in which F_1, F_0 are objects of $\text{Rep}(G)_{\text{flat}}$ and hence are described by our theorem. We conclude that E has the following form. There exist an

injective homomorphism $\phi : N \rightarrow N'$ of graded B -modules and flat A -submodules M, M' of N, N' respectively such that $\phi M \subset M'$ and $(N, M), (N', M') \in \mathcal{C}$, having the property that E is isomorphic to $M'/\phi M$ as G -module.

1.5. A special case. When A is a $\mathbb{Q}$ -algebra, the category $\mathcal{C}$ is very simple. Indeed, there is a polynomial $P_n \in \mathbb{Q}[U]$ of degree n such that $\binom{i}{n} = P_n(i)$, and therefore $C_n = Q_n(C)$, where $C = C_1$ and $Q_n := f^n P_n(f^{-1}U) \in A[U]$. Therefore $\mathcal{C}$ is the category of pairs (N, M) consisting of a $\mathbb{Z}$ -graded B -module N and an A -submodule M of N such that the natural map $B \otimes_A M \rightarrow N$ is an isomorphism and such that $CM \subset M$, where C is the endomorphism of the graded module $N = \bigoplus_{i \in \mathbb{Z}} N_i$ given by multiplication by fi on N_i .

When A is the formal power series ring $\mathcal{O} := \mathbb{C}[[\varepsilon]]$, and $f = \varepsilon^k$ (for some non-negative integer k) our constructions yield a group scheme G over $\mathcal{O}$ such that $G(\mathcal{O}) = \{t \in \mathcal{O}^\times : t \equiv 1 \pmod{\varepsilon^k}\}$, and the category of representations of G on free $\mathcal{O}$ -modules of finite rank is equivalent to the category of pairs (V, M) , where V is a finite dimensional graded vector space over $F := \mathbb{C}((\varepsilon))$ and M is an $\mathcal{O}$ -lattice in V such that $CM \subset M$, where C is given by multiplication by $i\varepsilon^k$ on the i -th graded piece of V . It is amusing to note that for fixed V , the space of all M satisfying $CM \subset M$ is an affine Springer fiber, which, when all the non-zero graded pieces of V are one-dimensional, is actually one of the affine Springer fibers studied at some length in [GKM], where it was shown to be paved by affine spaces. Finally, since $\mathcal{O}$ is a principal ideal domain, the results in 1.4 give a concrete description of all G -modules.

2. CERTAIN HOPF ALGEBRAS AND THEIR COMODULES

Throughout this section A is a commutative ring and B is a commutative algebra such that the canonical homomorphism $B \otimes_A B \rightarrow B$ (given by $b_1 \otimes b_2 \mapsto b_1 b_2$) is an isomorphism. For example B might be of the form $S^{-1}A/I$ for some multiplicative subset S of A and some ideal I in $S^{-1}A$.

Let N be a B -module. Then the canonical B -module map $B \otimes_A N \rightarrow N$ (given by $b \otimes n \mapsto bn$) is an isomorphism. It follows that the canonical A -module homomorphism $N \rightarrow B \otimes_A N$ (given by $n \mapsto 1 \otimes n$) is actually an isomorphism of B -modules (since $N \rightarrow B \otimes_A N \rightarrow N$ is the identity).

Moreover, for any two B -modules N_1, N_2 , we have isomorphisms

$$(2.0.1) \quad \text{Hom}_B(N_1, N_2) \cong \text{Hom}_A(N_1, N_2)$$

and

$$(2.0.2) \quad N_1 \otimes_A N_2 \cong N_1 \otimes_B N_2.$$

2.1. General remarks on Hopf algebras and their comodules. Let S be a Hopf algebra over A . The composition $A \rightarrow S \rightarrow A$ of the unit and counit is the identity, and therefore there is a direct sum decomposition $S = A \oplus S_0$ of A -modules, where S_0 is by definition the kernel of the counit $S \rightarrow A$. In this subsection all tensor products will be taken over A and the subscript A will be omitted.

We denote by $\Delta : S \rightarrow S \otimes S$ the comultiplication for S . The counit axioms imply that Δ takes the form $\Delta(a + s_0) = a + s_0 \otimes 1 + 1 \otimes s_0 + \bar{\Delta}(s_0)$, when we identify S with $A \oplus S_0$ and $S \otimes S$ with $A \oplus (S_0 \otimes A) \oplus (A \otimes S_0) \oplus (S_0 \otimes S_0)$. Here $\bar{\Delta}$ is a uniquely determined A -module map $S_0 \rightarrow S_0 \otimes S_0$.

For any S -comodule M with comultiplication $\Delta_M : M \rightarrow S \otimes M$ the counit axiom for M implies that $\Delta_M(m) = 1 \otimes m + \bar{\Delta}_M(m)$ for a uniquely determined A -module map

$$\bar{\Delta}_M : M \rightarrow S_0 \otimes M.$$

In this way we obtain an equivalence of categories between S -comodules and A -modules M equipped with an A -linear map $\bar{\Delta}_M : M \rightarrow S_0 \otimes M$ such that the diagram

$$(2.1.1) \quad \begin{array}{ccc} M & \xrightarrow{\bar{\Delta}_M} & S_0 \otimes M \\ \bar{\Delta}_M \downarrow & & \bar{\Delta} \otimes id \downarrow \\ S_0 \otimes M & \xrightarrow{id \otimes \bar{\Delta}_M} & S_0 \otimes S_0 \otimes M \end{array}$$

commutes.

2.2. Hopf algebras for B give Hopf algebras for A . Let S be a Hopf algebra over B . As in the previous subsection we decompose S as $B \oplus S_0$. It is easy to see that there is a unique Hopf algebra structure on $R := A \oplus S_0$ such that the unit and counit for R are the obvious maps $A \hookrightarrow R$ and $R \rightarrow A$ and such that the Hopf algebra structure on $B \otimes_A R$ agrees with the given one on S under the natural B -module isomorphism $B \otimes_A R \cong S$. What makes this work is (2.0.2), a consequence of our assumption that $B \otimes_A B \rightarrow B$ is an isomorphism, so that, for example, $S_0 \otimes_B S_0 \cong S_0 \otimes_A S_0$. The comultiplications Δ_R, Δ_S on R, S respectively are given by

$$(2.2.1) \quad \Delta_R(a + s_0) = a + s_0 \otimes 1 + 1 \otimes s_0 + \bar{\Delta}(s_0)$$

$$(2.2.2) \quad \Delta_S(b + s_0) = b + s_0 \otimes 1 + 1 \otimes s_0 + \bar{\Delta}(s_0)$$

and similar considerations apply to the multiplication maps $R \otimes_A R \rightarrow R$, $S \otimes_B S \rightarrow S$ and the antipodes $R \rightarrow R$, $S \rightarrow S$.

Proposition 2.2.1. *The category of R -comodules is equivalent to the category of A -modules M equipped with an S -comodule structure on $N := B \otimes_A M$.*

Proof. We have already observed that giving an R -comodule is the same as giving an A -module M equipped with an A -module map $\bar{\Delta}_M : M \rightarrow S_0 \otimes_A M$ such that (2.1.1) commutes. Since S_0 is a B -module and $B \otimes_A B \cong B$, giving $\bar{\Delta}_M$ such that (2.1.1) commutes is the same as giving a B -module map $\bar{\Delta}_N : N \rightarrow S_0 \otimes_B N$ such that

$$\begin{array}{ccc} N & \xrightarrow{\bar{\Delta}_N} & S_0 \otimes_B N \\ \bar{\Delta}_N \downarrow & & \bar{\Delta} \otimes id \downarrow \\ S_0 \otimes_B N & \xrightarrow{id \otimes \bar{\Delta}_N} & S_0 \otimes_B S_0 \otimes_B N \end{array}$$

commutes, or, in other words, giving an S -comodule structure on N . $\square$

2.3. Special case. Let $\mathcal{O}$ be a valuation ring and F its field of fractions. Let G be an affine group scheme over F and let S be the corresponding commutative Hopf algebra over F . Decompose S as $F \oplus S_0$ and define a commutative Hopf algebra R over $\mathcal{O}$ by $R := \mathcal{O} \oplus S_0$. Corresponding to R is an affine group scheme $\tilde{G}$ over $\mathcal{O}$, and giving a representation of $\tilde{G}$ (that is, an R -comodule) is the same as giving an $\mathcal{O}$ -module M together with an S -comodule structure on $F \otimes_{\mathcal{O}} M$.

For example, when G is the multiplicative group $\mathbb{G}_m$, the Hopf algebra R is $\{\sum_{i \in \mathbb{Z}} a_i T^i \in F[T, T^{-1}] : \sum_{i \in \mathbb{Z}} a_i \in \mathcal{O}\}$, which is easily seen to be the union of the Hopf subalgebras S_k discussed in the introduction.

REFERENCES

- [GKM] M. Goresky, R. Kottwitz and R. MacPherson, *Purity of equivalued affine Springer fibers*, Represent. Theory **10** (2006), 130–146.
- [Sa] N. Saavedra Rivano, *Catégories Tannakiennes*, Lecture Notes in Mathematics 265, Springer-Verlag, Berlin, 1972.
- [Se] J.-P. Serre, *Groupes de Grothendieck des schémas en groupes réductifs déployés*, Publ. Math. IHES **34** (1968), 37–52.
- [SGA3] M. Demazure and A. Grothendieck, *SGA3, Schémas en Groupes, Tome I*, Lecture Notes in Mathematics 151, Springer-Verlag, Heidelberg, 1970.
- [We] T. Wedhorn, *On Tannakian duality over valuation rings*, J. Algebra **282** (2004), 575–609.

N. E. CSIMA, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, 5734 UNIVERSITY AVENUE, CHICAGO, ILLINOIS 60637

E-mail address: `ecsima@math.uchicago.edu`

ROBERT E. KOTTWITZ, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, 5734 UNIVERSITY AVENUE, CHICAGO, ILLINOIS 60637

E-mail address: `kottwitz@math.uchicago.edu`